

Label

White papers

Coupons

Other

Image

MERCK ANIMAL HEALTH Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Banamine®



Intervet/Merck Animal Health

PRODUCT INFORMATION

(flunixin meglumine paste)

Paste - 1500 mg flunixin/syringe

Veterinary

For Oral Use in Horses Only

CAUTION Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION Each 30-gram syringe of BANAMINE Paste contains flunixin meglumine equivalent to 1500 mg flunixin.

INDICATIONS BANAMINE Paste is recommended for the alleviation of inflammation and pain associated with musculoskeletal disorders in the horse.

ACTIVITY Flunixin meglumine is a potent, nonnarcotic, nonsteroidal, analgesic agent with anti-inflammatory and antipyretic activity. It is significantly more potent than pentazocine, meperidine, and codeine as an analgesic in the rat yeast paw test. Oral studies in the horse show onset of flunixin activity occurs within 2 hours of administration. Peak response occurs between 12 and 16 hours and duration of activity is 24 to 36 hours.

CONTRAINDICATIONS There are no known contraindications to this drug when used as directed.

WARNING Not for use in horses intended for human consumption.

PRECAUTIONS The effect of BANAMINE Paste on pregnancy has not been determined. Studies to date show there is no detrimental effect on stallion spermatogenesis with or following the recommended dose of BANAMINE Paste.

SIDE EFFECTS During field studies with BANAMINE Paste, no significant side effects were reported.

DOSAGE AND ADMINISTRATION The recommended dose of flunixin is 0.5 mg per lb of body weight once daily. The BANAMINE Paste syringe, calibrated in twelve 250-lb weight increments, delivers 125 mg of flunixin for each 250 lbs (see dosage table). One syringe will treat a 1000-lb horse once daily for 3 days, or three 1000-lb horses one time.

DOSAGE TABLE			
Syringe Mark*	Horse Weight (lbs)	BANAMINE Paste Delivered (g)	Mg Flunixin Delivered
0	-	-	-
250	250	2.5	125
500	500	5.0	250
750	750	7.5	375
1000	1000	10.0	500

*Use dial edge nearest syringe barrel to mark dose.

The paste is orally administered by inserting the nozzle of the syringe through the interdental space, and depositing the required amount of paste on the back of the tongue by depressing the plunger.

Treatment may be given initially by intravenous or intramuscular injection of BANAMINE Solution, followed by BANAMINE Granules or BANAMINE Paste on Days 2 to 5. BANAMINE treatment should not exceed 5 consecutive days.

TOXICITY No toxic effects were observed in rats given oral flunixin 2 mg/kg per day for 42 days. Higher doses produced ulceration of the gastrointestinal tract. The emetic dose in dogs is between 150 and 250 mg/kg. Flunixin was well tolerated in monkeys dosed daily with 4 mg/kg for 56 days. No adverse effects occurred in horses dosed orally with 1.0 or 1.5 mg/lb for fifteen consecutive days.

HOW SUPPLIED BANAMINE Paste, 1500 mg is available in a single 30-g syringe.

Store below 25°C (77°F). Do not Freeze.

For patent information: <http://www.merck.com/product/patent/home.html>

NADA #137-409, Approved by FDA.

Made in France.

Copyright © 2011 Intervet International B.V, a subsidiary of Merck & Co., Inc., Madison, NJ, USA

All rights reserved

Net Wt	NDC	
30 g	0061-0214-02	160549 R1

CPN: 1047019.5

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

Copyright 2019 - Animalytix LLC